


ENDOSCOPY CENTRE

Preparing for a Colonoscopy:

For Patients Prescribed with PICOPREP (8AM TO 2PM)

For more
information on
Colonoscopy:



Dietary Restrictions Three Days Before Colonoscopy

Take only light, low-fibre meals such as:



White bread
(no spread)



White porridge/
rice



Lean meat
(chicken/fish)



Eggs



Coffee/tea (with
or without milk)



Clear soup



Mee Sua/Bee
hoon/Kway teow

Stop taking any oily and high-fibre foods such as:



Fried/oily
foods



Fruits



Vegetables



Whole
grains/ oats



Cereal

Medications for preparation: PICOPREP / PICOPREP and PEG

I. PICOPREP

- This is a drug (laxative) which stimulates the muscle in the intestines, in order to create bowel movements (pass motion).
- You may experience stomach cramps and diarrhoea after taking the medication.



Fig 1. One box of PICOPREP and cup provided

2. Polyethylene Glycol Electrolyte Solution (PEG)

- It is used to clean out the gastrointestinal tract (stomach and intestines).
- You may experience diarrhoea, nausea and/or stomach cramps after taking the medication.



Fig 2. Two packets of PEG powder

Important Notes:

- ✓ Do see a doctor at your nearest clinic if you experience vomiting or severe abdominal pain (stomachache).
- ✓ Depending on your procedure type, you will be required to take either:
 - "PICOPREP" (Page 4) OR
 - "PICOPREP + PEG" (Page 5 and 6)
- ✓ Please refer to the correct instructions for your procedure preparation

Medications for preparation: PICOPREP / PICOPREP and PEG

7 DAYS BEFORE Your Colonoscopy (Date:_____)

Stop and do not take these medications: _____.

- E.g. Iron, Plavix, Anti-diarrhoeal medications



Other comments:

(Not Applicable)

3 DAYS BEFORE Your Colonoscopy (Date:_____)

Low residue / low fibre diet (Refer to page I)

2 DAYS BEFORE Your Colonoscopy
(Date:_____)

Stop and do not take these medications: _____.

- E.g. Empagliflozin, dapagliflozin



(Not Applicable)



PICOPREP

(Follow instructions on Page 4)

1 DAY BEFORE Your Colonoscopy (Date: _____)

7AM Light low-fibre breakfast.

12PM Light low-fibre lunch.

2PM-3PM Prepare the PICOPREP solution to drink by mixing 1 packet of powder with 150 ml of warm water, and stirring for 2 to 3 minutes.



Drink at least 1 litre or 2 cups of plain water (1 cup = 500ml), using the PICOPREP cup given.

6PM-6.30PM Light low-fibre dinner.
No more food allowed after 6.30PM.

8PM-9PM Prepare the PICOPREP solution to drink by mixing 1 packet of powder with 150 ml of warm water, and stirring for 2 to 3 minutes.



Drink at least 1 litre or 2 cups of plain water (1 cup = 500ml), using the PICOPREP cup given.

ON THE DAY of Your Colonoscopy (Date: _____)

12 midnight onwards No food allowed

Time _____
(2 hours before the procedure)

- Stop and do not take these medications _____.
- Continue other usual medications. Take it at least 2 hours before your colonoscopy, with small amounts of water.
- Stop drinking fluids including plain water

Time _____ Report to Endoscopy Centre



PICOPREP + PEG

(Follow instructions on Page 5 and 6)

I DAY BEFORE Your Colonoscopy (Date:_____)

7AM Light low-fibre breakfast.

12PM Light low-fibre lunch.

2PM-3PM Prepare the PICOPREP solution to drink by mixing 1 packet of powder with 150 ml of warm water and stirring for 2 to 3 minutes.

Drink at least 1 litre or 2 cups of plain water (1 cup = 500ml), using the PICOPREP cup given.

6PM-7PM Light low-fibre dinner.
No more food allowed after 7PM.

8PM-9PM Prepare the PICOPREP solution to drink by mixing 1 packet of powder with 150 ml of warm water and stirring for 2 to 3 minutes.

Drink at least 1 litre or 2 cups of plain water (1 cup = 500ml), using the PICOPREP cup given.





PICOPREP + PEG

(Follow instructions on Page 5 and 6)

ON THE DAY of Your Colonoscopy (Date: _____)

12 midnight
onwards

No food allowed

5AM-6AM

- Stop and do not take these medications_____.
- Continue other usual medications. Take it at least 2 hours before your colonoscopy, with small amounts of water.



Mix 1 packet of PEG powder with 1 litre or 4 cups of plain water (1 cup = 250 ml). Start drinking 1 litre of PEG solution from 5AM and complete by 6AM.



- Stop drinking fluids including plain water
- No food allowed

Time _____: Report to Endoscopy Centre

Good vs. Poor Bowel Preparation

✗ Poor Bowel Preparation Leads to a Dirty Colon

Please report to the Endoscopy Centre 2 hours before your appointment time if your stool appears like any of the pictures shown below (Stool 1 to 4).


Your doctor may advise additional medications for your bowel preparation.

Stool Number Chart:



✓ Good Bowel Preparation Leads to a Clean Colon

Stool Number Chart:



Endoscopy Pre-Admission Counselling
and Evaluation (Endo-PACE)

TTSH Medical Centre, Level 2
Contact:
6357 7000 (Central Hotline)



Scan the QR Code with your smart phone
to access the information online or visit
<https://nhghealth.com.sg>



© Tan Tock Seng Hospital, Singapore 2026. All rights reserved. All information correct as of March 2026. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.